arXiv:1601.06278v1 [cs.DM] 23 Jan 2016

Fortuitous sequences of flips of the top of a stack of n burnt pancakes for all $n > 24$.

Laurent PIERRE, université de Nanterre, France
Laurent.PIERRE@u-paris10.fr

Summary

Burnt pancakes problem was defined by Gates and Papadimitriou in 1979. They proved that $g(-f_n) \geq 3n/2 - 1$.

In 1995 Cohen and Blum conjectured that $g(n) = g(-I_n)$ and proved that $g(-I_n) = g(-f_n) + 1 \geq 3n/2$.

In 1997 Heydari and Sudborough proved that $g(-I_n) \leq 3\frac{n+1}{2}$ whenever some "fortuitous sequence" exists. They gave fortuitous sequences for $n = 3$, 15, 27 and 31. They showed that two fortuitous sequences S_n and S'_n may combine into another fortuitous sequence S''_n with $n'' = n + n' - 3$. So a fortuitous sequence S_n gives a fortuitous sequence S_{n+12}. This proves that $g(-I_n) \leq 3\frac{n+1}{2}$ if $n \equiv 3 \pmod 4$ and $n \geq 23$.

In 2011 Josef Cibulka enhanced Gates and Papadimitriou's lower bound thanks to a potential function. He got so $g(-I_n) \geq 3n/2 + 1$ if $n > 1$ proving thereby, that $g(-I_n) = 3\frac{n+1}{2}$ if $n \equiv 3 \pmod 4$ and $n \geq 23$. Cibulka gave also a sequence of 30 flips (19, 14, 7, 4, 10, 18, 6, 4, 10, 19, 14, 4, 9, 11, 8, 18, 8, 11, 9, 4, 14, 19, 10, 4, 6, 18, 10, 4, 7, 14) sorting $-I_{19}$, without being fortuitous. With the fortuitous sequence for 15 pancakes this proved that $g(-I_n) = 3\frac{n+1}{2}$ if $n \equiv 3 \pmod 4$ and $n \geq 15$. Cibulka also invalidated Cohen and Blum's conjecture by showing that $g(-I_{15}) = 24$ while $g(15) = 25$.

This paper explains how to build generalized fortuitous sequences for every $n \geq 25$ proving thereby that $g(-I_n) = \lceil \frac{3}{2}n \rceil + 1$ for every $n \geq 25$.

Burnt pancakes problem

Stack of burnt pancakes $\begin{bmatrix} 2 \\ -4 \\ 5 \\ 1 \\ -3 \end{bmatrix}$ holds 5 pancakes numbered from 1 to 5 from smallest to largest. In this stack pancakes 3 and 4 are upside down. In order to save space, it will be rather written S=[2 -4 5 1 -3], top of stack ahead.

This stack of pancakes may be seen as a signed permutation of integers 1 to 5, i.e. mapping $1 \mapsto 2$, $2 \mapsto -4$, $3 \mapsto 5$, $4 \mapsto 1$, $5 \mapsto -3$. This mapping may be extended to negative integers -5 to -1 by $S(-i) = -S(i)$. Such mappings may be composed. E.g. [2 -4 5 1 -3]$\circ$[1 2 -3 5 4]=[2 -4 -5 -3 1].

Stack I_n=[1 2 3 ... n] denotes identity mapping. It is also target setting of stack of pancakes problem. Pancakes are sorted by size, smallest one on top. They are all well oriented i.e. burnt side down.

Stack $-I_n$=[-1 -2 -3 ... -n] is initial setting of stack of pancakes problem. Pancakes are all in right location but they are upside down, i.e. burnt side up.

Spatula flip f_i splits stack of pancakes in two parts by inserting a spatula below the i topmost pancakes, and then flips upper part. E.g. f_3=[-3 -2 -1 4 5] when stack holds 5 pancakes. Flips are only operations allowed to sort stack of pancakes.

Any signed permutation S of a stack of n burnt pancakes, may be achieved by a flip sequence. Let $g(S)$ be the minimal number of flips needed. Cibulka proved in 2011 that $g(-I_n) \geq 3n/2 + 1$ if $n > 1$.

Flip sequence $w = (f_4\ f_8\ f_6\ f_{10})$ will be shortened as $w = (4\ 8\ 6\ 10)$. Let $\tilde{w} = (10\ 6\ 8\ 4)$ and $f_w = f_{10} \circ f_6 \circ f_8 \circ f_4$. Permutation f_w is effect of sequence w. Applying flip sequence S to stack $-I_n$ gives stack $f_S \circ -I_n = -f_S$. So "sequence S sorts stack $-I_n$" translates into $f_S \circ -I_n = I_n$ i.e. $f_S = -I_n$.

Lower bound on $g(-I_n)$ by Cohen and Blum : $3n/2$

When Gates and Papadimitriou introduced burnt pancake problem in 1979, they demanded also that pancakes in initial setting had right orientation. In other word every single pancake had to be flipped an even number of times. That is why, they looked at $g(-f_n)$ instead of $g(-I_n)$: at start all pancakes have right orientation, but they are sorted in reversed order $-f_n$=[n n-1 ... 3 2 1]. They proved that $g(-f_n) \geq 3n/2 - 1$.

In 1995 Cohen and Blum noticed that every signed permutation p of a stack of n pancakes verifies $p \circ -I_n = -I_n \circ p = -p$. Hence $-I_n$ commutes to every signed permutation. (In other word $-I_n$ is in center of group of signed permutations. Indeed center is $\{I_n, -I_n\}$).

If $-I_n = p \circ q$ then $-I_n$ commutes to p and p^{-1} and so $-I_n = q \circ p$. This means that a flip sequence which sorts $-I_n$, may be permuted circularly. It will still sort $-I_n$. Similarly $-I_n$ and flips are their own inverses, hence a flip sequence sorting $-I_n$ may be read backward, it will still sort $-I_n$. For $n > 1$, a flip sequence sorting $-I_n$ holds at least two f_n, first one removes -n from bottom of stack, while last one brings n back at bottom. Hence minimal sequences sorting $-I_n$ may be permuted circularly to produce other minimal sequences starting by f_n. Removing this initial f_n produces a minimal sequence sorting $-f_n$. Hence $g(-I_n) = 1 + g(-f_n)$. This is how Cohen and Blum proved that $g(-I_n) \geq 3n/2$.

Lower bound for $g(-I_n)$ with a potential function

In a stack of pancakes a "block" is a maximal sequence of several contiguous pancakes which appears also in I_n or f_n. A "clan" is a maximal sequence of several contiguous pancakes which appears also in $-I_n$ or $-f_n$. Two contiguous pancakes in a same block are "adjacent". Two contiguous pancakes in a same clan are "anti-adjacent". A pancake which is neither in block nor in a clan is a "singleton". E.g. stack [4 5 6 -1 -2 3] is made of block (4 5 6), clan (-1 -2) and singleton 3. Potential of a pancake in a stack will be defined as 1 when at an end of a block, 3 when inside a block, -1 when at an end of a clan, -3 when inside a clan and 0 for a singleton. Potential of top pancake is doubled. Potential of a stack is sum of potentials of its pancakes. E.g potential of stack $\begin{matrix}[4\ 5\ 6\ {-1}\ {-2}\ 3] \\ 2\ 3\ 1\ {-1}\ {-1}\ 0\end{matrix}$ is 4.

During flip $[-x \dots y|z \dots] \rightarrow [-y \dots x|z \dots]$ potential function varies only for pancakes x, y and z. Possible variations of $p(y)$ are $3 \rightarrow 2$, $1 \rightarrow 2$, $1 \rightarrow 0$, $0 \rightarrow 0$, $-1 \rightarrow 0$, $-1 \rightarrow -2$ and $-3 \rightarrow -2$. Hence $|\Delta p(y)| \leq 1$. As well $|\Delta p(x)| \leq 1$. Possibles changes of $p(z)$ are $3 \leftrightarrow 1$, $1 \leftrightarrow 0$, $0 \leftrightarrow -1$, $1 \leftrightarrow -1$ and $-1 \leftrightarrow -3$. Hence $|\Delta p(z)| \leq 2$. Hence $|\Delta p(x) + \Delta p(y) + \Delta p(z)| \leq 1 + 1 + 2 = 4$.

During flip $[-x|z \dots] \rightarrow [x|z \dots]$ potential function varies only for pancakes x and z. Possible variations of $p(x)$ are $2 \leftrightarrow 0$, $0 \rightarrow 0$ and $-2 \leftrightarrow 0$. Hence $|\Delta p(x)| \leq 2$. We still have $|\Delta p(z)| \leq 2$. Hence $|\Delta p(x) + \Delta p(z)| \leq 4$.

This proves that during a flip potential function increases by at most 4.

$p(I_n) = 2 + 3 + 3 + \dots + 3 + 1 = 3n - 3$. $\quad p(-I_n) = p(-f_n) = -p(I_n) = 3 - 3n$.

Hence $\quad g(-f_n) \geq \dfrac{p(I_n) - p(-f_n)}{4} = \dfrac{3n-3}{2} \quad$ and $\quad g(-I_n) \geq \dfrac{3n-1}{2}$.

Enhancement of potential function to reach Cibulka's bound : $g(-I_n) \geq 3n/2 + 1$

Potential function may be enhanced if plate supporting the stack behaves like a pancake, which may form a block with bottom pancake when it is n or a clan if it is 1. Plate potential is -1, 0 or 1 and adds to potential function of stack. Final potential is now $p(I_n) = 3n$ and initial potential is $p(-f_n) = -3n$. Then $g(-f_n) \geq 6n/4 = 3n/2$. Hence $g(-I_n) \geq 3n/2 + 1$.

```
[ 3 2 1]      [-2-3 1]      [-1 3 2]      [-3 1 2]      [-2-1 3]       [1 2 3]
 -2-3-3-1      -2-1-1-1       0-1-1         0 1 1         2 1 1 1       2 3 3 1
   -9            -5             -2            2             5             9
         4             3             4             3             4
```

Since $g(-I_n)$ is an integer, the lower bound is actually : $g(-I_n) \geq \left\lceil \frac{3n}{2} + 1 \right\rceil = \left\lfloor \frac{3n+3}{2} \right\rfloor$.

This was proved in 2011 by Cibulka who used about the same potential function with the following differences :

* His potential function is the third of ours. E.g. potential function variation during a flip is at most 4/3.

* Potential of a block with n_a adjacencies is $3n_a$ if it is on top of stack and $3n_a - 1$ if it is inside stack. As well potential of a clan with n'_a anti-adjacencies is $-3n'_a$ when in top of stack and $-3n'_a + 1$ otherwise. Hence potential of a stack S is $p(S) = p'(S) - p'(-S)$ with $p'(S) = 3n_a(S) - n_b(S)$ where n_a and n_b count adjacencies and inner blocks. Cibulka takes $p(S) = n_a(S) - n_a(-S) - (n_b(S) - n_b(-S))/3$.

* Cibulka keeps a neutral plate under the stack of pancakes, but he adds to potential 4/3, 1, -1 or -4/3 if bottom of stack is n-1,n or n or -n or 1-n,-n respectively.

* Cibulka starts from $-I_n$ instead of $-f_n$, but he adds 1/3 to potential when a block holds 1,2 (or -2,-1) or when -1 is on top of stack. As well he adds -1/3 to potential when a clan holds 2,1 (or -1,-2) or when 1 is on top of stack.

Flip sequence reaching lower bound

Let us assume $g(-I_n) = \lceil 3n/2 + 1 \rceil$. If n is even, then $g(-f_n) = 3n/2$. There exists a flip sequence of length $3n/2$ transforming $-f_n$ into I_n. Potential function goes from $-3n$ to $3n$ in $3n/2$ flips. Hence every flip increases potential by 4. For every flip f_1 we have $\Delta p(x) = \Delta p(z) = 2$. For every larger flip we have $\Delta p(x) = \Delta p(y) = 1$ and $\Delta p(z) = 2$. Potential function of a pancake increases by 1 when it leaves or reaches top of stack, and increases by 2 when not moved. This proves that parity of potential of a pancake depends only on its location in stack : Every pancake but top one has odd potential. Only top pancake may be a singleton.

Plate potential starts from -1 in $-f_n$ and ends at 1 in I_n. It increases by 2 during every f_n. So only one f_n is used to transform $-f_n$ into I_n, and two of them are used to transform $-I_n$ into I_n.

If n is odd, then lower bound is reached when $g(-I_n) = (3n + 3)/2$. There exists a flip sequence of length $(3n + 1)/2$ transforming $-f_n$ into I_n. Potential function goes from $-3n$ to $3n$ in $(3n + 1)/2$ flips. Potential variation is loose by 2. There may be two f_n. Then first f_n changes plate potential from -1 to 0 and second f_n from 0 to 1. Variation of potential during both f_n will be 3. Hence variation of potential for every smaller flip will be 4. Transformation from $-I_n$ to I_n uses three f_n.

We shall build flip sequences reaching lower bound for every n large enough.

Fortuitous sequences and their combinations

A "fortuitous sequence" is a flip sequence of the form $S = ns_1ns_2ns_3$ where s_1, s_2 and s_3 are sequences of length $\frac{n-1}{2}$ of even positive integers lower than n, such that $f_S = -I_n$.

So n is odd and $|S| = \frac{3n+3}{2}$. Lower bound is reached. Hence when applying S to $-I_n$, sequence $s_1ns_2ns_3$ applies do $-f_n$ to reach I_n and every flip increases potential function by 4, but both f_n which increase potential by only 3. Since s_1 is made of even numbers, when applying s_1 to $-f_n$, pancake 1 stays at bottom of stack, and above it is a stack of clans of even lengths. Each flip, to increase potential function by 4, must break a clan into two smaller clans. At the end of s_1 the stack $f = -f_{ns_1}$ is a stack of $\frac{n-1}{2}$ clans of 2 pancakes placed on singleton 1. Heydari and Sudborough called it the first column of a box. We shall call it a "stack of 2-clans". Similarly every flip of s_3 glues two blocks to form a bigger one. And stack $-f_{ns_1ns_2n}$ at the beginning of s_3 is a stack of $\frac{n-1}{2}$ blocks of 2 pancakes, placed on singleton n. Hence every flip of s_2 replaces a 2-clan by a 2-block. First flip of s_2 is $[-1 \dots 3|2 \dots] \to [-3 \dots 1|2 \dots]$. Second flip of s_2 is $[-3 \dots 5|4 \dots] \to [-5 \dots 3|4 \dots]$. Third flip of s_2 is $[-5 \dots 7|6 \dots] \to [-7 \dots 5|6 \dots]$. Etc.. So during application of flip sequence s_2 top pancakes becomes successively -1, -3, -5, -7, ..., $-n$. Bottom pancake does not move, but it leaves its initial clan to join its final block during the only f_{n-1}. This proves that s_2 holds exactly one occurrence of $n-1$ and that bottom pancake is even and positive i.e. right-side up. So top pancake of stack of 2-clans f is even and negative i.e. upside down. Of course s_1 holds also exactly one occurrence of $n-1$, and s_3 too.

Heydari et al. demanded that top pancake in the stack of 2-clans should be $-\frac{n+1}{2}$, but we do not. For us, it is only even and negative.

We may notice that $h = f_n \circ f^{-1} \circ f_n$ is the stack of 2-clans of fortuitous sequence $n\tilde{s}_1n\tilde{s}_3n\tilde{s}_2$ and that s_1, $\tilde{s}_1$, s_2 and $\tilde{s}_3$ are the only flip sequences adding an adjacency at each step when starting respectively from stacks $-h$ (or $-f_n \circ f^{-1}$), $-f$, h^{-1} (or $f_n \circ f$) and f^{-1} (or $f_n \circ h$). So stack of 2-clans f characterizes flip sequence S.

We may roughly count fortuitous sequences. First flip of s_1 is a number taken among 2, 4, 6, ... n-1. There are $\frac{n-1}{2}$ possibilities. Second flip must break a clan. There are $\frac{n-1}{2} - 1$ possibilities. Third flip is to be chosen among $\frac{n-1}{2} - 2$ possibilities. Etc.. So there are $\frac{n-1}{2}!$ ways of choosing s_1 to get a stack of 2-clans. After that s_2 and s_3 are imposed. Indeed there is only one way to create an adjacency by a flip. This flip has about one chance out of two to exist, whether pancake matching first one shows right side or not. So expected number of fortuitous sequences may be estimated as $\frac{n-1}{2}!/2^{n-1} \sim \sqrt{\pi n}\left(\frac{n-1}{8e}\right)^{\frac{n-1}{2}} \to \infty$. This number is large as soon as $n > 8e + 3$. This suggests, that fortuitous sequences exist for every large enough

odd integer n. This can be proved in two different ways, that we shall both use. Either we look for regular patterns in stacks of 2-clans of small fortuitous sequences and design according to these patterns a stack of 2-clans yielding a fortuitous sequence for any large odd integer n. Or we exhibit small fortuitous sequences with symmetries, like palindromic or triple fortuitous sequences and use method of Heidari et al. to combine two fortuitous sequences for n_1 and n_2 pancakes into a fortuitous sequence for $n_1 + n_2 - 3$ pancakes. E.g. fortuitous sequences
(15 10 4 6 14 6 4 10 15 10 4 6 14 6 4 10 15 10 4 6 14 6 4 10)
(23 14 4 6 22 10 8 12 10 14 12 18 23 10 14 18 8 10 22 10 8 18 14 10 23 18 12 14 10 12 8 10 22 6 4 14)
yield stacks of 2-clans :
[-8 -9 -12 -13 -2 -3 -14 -15 5 4 7 6 -10 -11|1]
[-12 -13|-16 -17 -6 -7 -18 -19 -4 -5 -20 -21 -2 -3 -22 -23 9 8 11 10 -14 -15 1]
In second stack first 2-clan -12(=-x) -13 is replaced by first stack without final 1. Also in first stack pancakes 2... 15(=n_1) are renumbered 12(=x)... 25 by increasing their numbers by 10(=$x-2$). Similarly in second stack pancakes 14(=$x+2$)... 23(=n_2) are renumbered 26... 35(=n_1+n_2-3) by increasing their numbers by 12(=n_1-3). This yields stack of 2-clans
[-18 -19 -22 -23 -12 -13 -24 -25 15 14 17 16 -20-21|-28-29 -6 -7-30-31 -4 -5-32-33 -2 -3-34-35 9 8 11 10-26-27 1]
and fortuitous sequence
(35 26 16 10 4 6 18 34 10 8 12 10 14 12 30 6 4 10 35 10 14 18 8 10 30 4 6 34 26 24 30 16 8 24 20 16 35 26 20 22 30 6 4 26 12 14 10 12 8 10 34 18 16 26)

Since there is a fortuitous sequence for 15 pancakes, any fortuitous sequence for n pancakes yields fortuitous sequences for $n+12$, $n+24$, $n+36$, etc.. Moreover a fortuitous sequence for n pancakes with gcd(n−3,12)=2 will yield fortuitous sequences for m pancakes for all large enough odd integer m. Indeed fortuitous sequences are known for 3, 15, 23, 29, 31, 33 and 37. For small n they can be found by trying all $\frac{n-1}{2}!$ possibilities. (6h on a PC for n=29). For large n search process must be improved. Fortuitous sequence $S = ns_1ns_2ns_3$ yields fortuitous sequences $ns_2ns_3ns_1$, $ns_3ns_1ns_2$, $n\tilde{s}_3n\tilde{s}_2n\tilde{s}_1$, $n\tilde{s}_2n\tilde{s}_1n\tilde{s}_3$ and $n\tilde{s}_1n\tilde{s}_3n\tilde{s}_2$. But these 6 sequences are not necessarily different. There are only 3 different sequences if $s_1 = \tilde{s}_1$, only 2 if $s_1 = s_2$ and only 1 if $s_1 = s_2 = \tilde{s}_1$. Indeed for n=3 and n=15 there is only one fortuitous sequence. Hence it verifies $s_1 = s_2 = \tilde{s}_1$. So it makes sense to search only for fortuitous sequences verifying $s_1 = \tilde{s}_1$ or $s_1 = s_2$.

Palindromic fortuitous sequences

When $s_1 = \tilde{s}_1$, then s_1 is a palindrome containing exactly one occurrence of $n-1$. So $s_1 = w\ n{-}1\ \tilde{w}$ is of odd length $\frac{n-1}{2}$ and $n \equiv 3 \pmod 4$. An exhaustive search of all $s_1 = w\ n{-}1\ \tilde{w}$ tries $\frac{n-1}{2}!/\frac{n+1}{4}!$ cases and finds readily all fortuitous sequences for n=27 (0.16s), n=31 (3.9s) and n=35 (2m 2s). E.g. it finds sequences of the form ($n\ w_n\ n{-}1\ \tilde{w}_n$) with
w_{23} = (14 4 6 22 10 8 12 10 14 12 18 23 10 14 18 8 10)
w_{27} = (22 16 10 18 26 6 4 12 14 22 6 4 10 27 12 4 8 22 4 6)
w_{31} = (18 4 14 22 6 4 18 26 6 4 22 30 6 4 26 31 10 4 18 4 26 4 14)
w_{35} = (6 22 18 10 32 8 26 22 34 22 26 8 32 10 18 22 6 35 6 22 18 10 32 8 26 22)

We may notice that if $f_{nwx\tilde{w}} = -I_n$ then $f_x = f_w^{-1} \circ -f_n \circ f_w$ and $-f_n$ are conjugate and have as many fixed points. If n is odd, $-f_n$ has one fixed point $\frac{n+1}{2}$. So f_x has one fixed point and $x = n-1$. If n is even, $-f_n$ has no fixed point like f_x. So $x = n$. So central flip of a palindromic flip sequence sorting $-f_n$ is always either f_{n-1} or f_n depending on parity of n, even if it is not a part of a fortuitous sequence.

Triple fortuitous sequences

For $n \equiv 1 \pmod 4$ we cannot have $s_1 = \tilde{s}_1$, but we may have $s_1 = s_2 = s_3$. Then the stack of 2-clans is a signed permutation of order 3. It is possible to search for all stacks of 2-clans of order 3, and keep among them those corresponding to fortuitous sequences. Let $d(i) = \mathrm{sgn}(i)(-1)^i$. A stack of 2-clans of order 3 is a permutation of $\{-n, 1{-}n, \ldots -1, 0, 1, 2, \ldots n\}$ verifying following conditions:

$*\forall i,\ f(-i) = -f(i)$	f is a signed permutation.
$*f(n) = 1$	Smallest pancake is at bottom right-side up.
$*f(1) < 0$	Top pancake is upside down.
$*\forall i,\ f(f(f(i))) = i$	f is a permutation of order 3.
$*\forall i,\ d(f(i)) = d(i)$	Orientation of a pancake depends on parities of its number and location.

$*\forall i,\ f(i-d(i)) = f(i)+d(i)$ f is made of clans of 2 pancakes.
$*\forall i \neq 0,\ f(i+d(i)) \neq f(i)-d(i)$ Clans of 2 pancakes do not form larger clans.

Depth first search is effective : We pick up any i such that $j = f(i)$ is known and $f(j)$ is unknown. Then we shall try all possible values k for $f(j)$. Absolute value of k must be taken in set $K = \{x{>}0 \mid f(x)$ and $f^{-1}(x)$ are unknown$\}$. Sign of k is determined by rule $d(k) = d(i)$, so $k = |k|d(|k|)d(i)$. Then we force $f(j) = k$ and $f(k) = i$. Furthermore whenever we force $f(x) = y$, we shall also force $f(-x) = -y$ and $f(x-d(x)) = y+d(y)$ and if only one of the numbers $f(y)$ and $f^{-1}(x)$ is known, then both numbers are forced to same value. First choice of $|k|$ is made in set $K = \{2, 3, \ldots, n-1\}$ since $f(n) = 1$. After each choice of k set K loses elements $|k|$, $|k+1|$, $|k-1|$, $|j-d(j)|$ and $|i+d(i)|$. Generally it loses at least 5 elements, since if for instance $|k+d(k)|$ was already excluded because $z = f(k+d(k))$ is known, then we shall force $f(z) = j-d(j)$ and exclude other elements. An exhaustive search of all stack of 2-clans of order 3 tries about $(n-2)(n-7)(n-12)(n-17)\ldots$ cases and finds readily all triple fortuitous sequences for n=29 (0.08s), n=31 (0.23s), n=33 (0.79s), n=35 (2.8s) and n=37 (10.5s). E.g.
(29 16 4 26 6 20 4 16 24 6 14 28 12 4 24)^3 (31 18 8 28 6 18 4 14 8 4 22 6 24 30 4 16)^3
(33 18 8 30 8 2 20 4 14 8 4 24 8 26 32 4 18)^3 (35 20 8 2 32 8 22 4 18 2 28 2 8 16 34 16 4 30)^3
(37 22 12 2 34 8 20 4 16 2 10 4 26 2 8 28 36 6 20)^3
Adding rule $f = f_n \circ f^{-1} \circ f_n$ gives readily all palindromic triple fortuitous sequences for n=35, 39, 43 and 47(0.57s). E.g. it gives (n w_n $n{-}1$ $\tilde{w}_n$)^3 with w_{35} =(6 22 18 10 32 8 26 22), w_{39} =(22 16 18 14 16 30 34 12 14), w_{43} =(6 26 22 14 40 10 12 10 30 26) and w_{47} =(6 28 24 16 44 12 2 14 10 32 28).

Even fortuitous sequences and their combination

For even n a sequence satisfying lower bound is of length $3n/2+1$ and holds two f_n. So we define an "even fortuitous sequence" as a flip sequence of length $3n/2+1$ of the form $S = ns_1ns_2$ where s_1 and s_2 are sequences of positive integers lower than n, such that $f_S = -I_n$ and stack $f = -f_{ns_1}$ obtained at the end of s_1, when applying S from $-I_n$ or equivalently s_1 from $-f_n$, is a "patchwork" of $\frac{n}{2}-1$ clans or blocks of 2 pancakes, placed on pancake 1 and below singleton $-n$.

So no pancake of this patchwork has potential 3 (or -3) and every flip of s_1 increases potential by 4, and breaks an anti-adjacency. As well every flip of s_2 creates an adjacency.

We may notice that $-f^{-1}$ and $f_n \circ f^{-1} \circ f_n$ and $-f_n \circ f \circ f_n$ are the patchworks of fortuitous sequences ns_2ns_1 and $n\tilde{s}_1n\tilde{s}_2$ and $n\tilde{s}_2n\tilde{s}_1$ respectively. Hence $\tilde{s}_1$, $\tilde{s}_2$, s_1, and s_2 are the only flip sequences adding an adjacency at each step when starting respectively from stacks $-f$, f^{-1}, $-f_n \circ f^{-1}$ and $f_n \circ f$. So patchwork f characterizes even fortuitous flip sequence S.

We may roughly count even fortuitous sequences. Number of patchworks is $\frac{n-2}{2}!2^{n-2}$, since there are $\frac{n-2}{2}!$ possible orders for the $\frac{n-2}{2}$ pieces of 2 pancakes, which have each 4 possible orientations (E.g. [2,3], [-2,-3], [3,2] or [-3,-2]). About a quarter of these patchworks are not suitable since they have too large blocks or clans. When choosing a patchwork, s_1 and s_2 are imposed. Indeed there is only one way to create an adjacency by a flip. But this flip has about one chance out of two to exist, whether pancake matching first one shows right side or not. Hence probability that s_2 exists is about $2^{-|s_2|}$ where $|s_2|$ the length of s_2 is the number of missing adjacencies in patchwork. As well probability that s_1 exists is about $2^{-|s_1|}$ where $|s_1|$ is number of missing anti-adjacencies in patchwork. Obviously $|s_1|+|s_2| = \frac{3n}{2}-1$, so expected number of fortuitous sequences may be estimated as $\frac{3}{4}\frac{n-2}{2}!2^{-1-n/2} \to \infty$. This suggests, that fortuitous sequences exist for every large enough even integer n. This can be proved, like for odd integers, in two different ways, that we shall both use. Either we look for regular patterns in patchworks of small even fortuitous sequences and design according to these patterns a patchwork yielding a fortuitous sequence for any large even n. Or we exhibit small even fortuitous sequences with symmetries, like palindromic or double fortuitous sequences and use method of Heidari et al. to combine two even fortuitous sequences for n_1 and n_2 pancakes into a fortuitous sequence for n_1+n_2-2 pancakes. E.g. even fortuitous sequences
(26 20 14 16 11 3 24 11 16 8 19 7 13 11 25 8 21 18 3 15 26 15 3 18 5 21 8 25 13 5 19 8 16 11 24 3 11 14 10 18)
(28 22 2 16 18 13 3 26 11 18 2 10 21 7 15 13 27 8 23 20 3 17 28 15 3 18 5 21 8 27 15 5 21 10 2 18 11 26 3 13 16 10 20) yield patchworks:
[-26 6 7 -21 -20 -23 -22 -16 -17 -3 -2 -13 -12 15 14 25 24 10 11 5 4 9 8 -18 -19|1]
[-28|6 7 -9 -8 -23 -22 -25 -24 -18 -19 -3 -2 -15 -14 17 16 27 26 12 13 5 4 11 10 -20 -21 1]
In second stack top singleton -28(=-n_2) is replaced by first stack without final 1. Also in first stack pancakes

2... 26(=n_1) are renumbered 28(=n_2)... 52(=n_1+n_2-2) by increasing their numbers by 26(=n_2-2). In second stack pancakes keep their numbers 1... 27(=n_2-1). This yields patchwork
[-52 32 33-47-46-49-48-42-43-29-28-39-38 41 40 51 50 36 37 31 30 35 34-44-45|6 7 -9 -8-23-22-25-24-18-19 -3 -2-15-14 17 16 27 26 12 13 5 4 11 10-20-21 1]
and even fortuitous sequence
(52 20 14 16 11 3 24 11 16 8 19 7 13 11 46 2 16 18 13 3 50 35 42 2 10 45 7 39 37 51 8 47 44 3 41 8 21 18 3 15 52 15 3 18 5 21 8 41 3 44 31 47 8 51 39 5 45 8 42 37 50 3 11 14 10 44 15 5 21 10 2 18 11 26 3 13 16 10 20)

Since there is a fortuitous sequence for 26 pancakes, any fortuitous sequence for n pancakes yields fortuitous sequences for $n+24$, $n+48$, $n+72$, etc.. Moreover even fortuitous sequences for 26 and 28 pancakes yield even fortuitous sequences for all n=$24i$+$26j$+2, i.e. for all large enough even integer n. Indeed fortuitous sequences are known for 2 and all even integers from 26 to 48. This yield fortuitous sequences for all even $n \geq 26$.

Searching even fortuitous sequences

Patchworks are permutations of $\{-n, 1-n, \ldots -1, 0, 1, 2, \ldots n\}$ verifying following conditions:

$*\forall i,\ f(-i) = -f(i)$	f is a signed permutation.
$*f(n) = 1$	Smallest pancake is at bottom right-side up.
$*f(1) = -n$	Largest pancake is on top up side down.
$*\forall i,\ f(i+d(i)) = f(i) + d(f(i))$	f is made of clans and blocks of 2 pancakes.
$*\forall i,\ 2f(i) \neq f(i-1) + f(i+1)$	Clans and blocks of 2 pancakes do not form larger clans or blocks.

E.g. $f(4) = -11$ implies $f(-4) = 11$, $f(5) = -10$, $f(-5) = 10$, $f(6) \neq -9$ and $f(3) \neq -12$. We perform a depth first search to find all even fortuitous sequences. At start patchwork f and stacks $-f$, f^{-1}, $-f_n \circ f^{-1}$ and $f_n \circ f$ have most of their values unknown. To make sure a flip on one of the four stacks creates an adjacency, we may have to define some values of f. We choose the stack which minimizes the number of unknown values of f to define. E.g. a stack of the form $[x \ldots y \ldots]$ is best choice, if values x and $y = 1-x$ are known, while a stack with an unknown top pancake is worst choice. Once the stack is chosen we try any possible flip on it and choose again a stack. When the top of a stack is 1 or $-n$ we may force the stack to be I_n or f_n respectively.

This depth first search finds even fortuitous sequences for $n = 26$(in 1m 23s) and $n = 28$(in 31m). Indeed it finds all solutions in 20m and 6h respectively, but for 26 as well as for 28, there are only 4 solutions, which yield one another : ns_1ns_2, ns_2ns_1, $n\tilde{s}_1n\tilde{s}_2$ and $n\tilde{s}_2n\tilde{s}_1$. We may look for similarities between a solution for 26 and a solution for 28. So we force $f(2) = 11$, $f(-4) = 21$, $f(-6) = 23$, $f(-8) = 2$, $f(-10) = 18$, $f(1-n) = 16$, $f(n-3) = 6$ and $f(n-5) = n-1$. This gives readily solutions for 30, 32, 34 and 38. Then looking for other similarities between those solutions, we impose
f = [$-n$ 11 10 -21 -20 -23 -22 -2 -3 -18 -19 w_n 13 12 -15 -14 8 9 5 4 w'_n n−2 n−1 7 6 -17 -16 1]
where w_n is an initial segment of length multiple of 4, of infinite sequence : -26 -27 -24 -25|-30 -31 -28 -29|-34 -35 -32 -33|-38 -39 -36 -37|-42 -43 -40 -41|-46 -47 -44 -45|-50 -51 -48 -49|-54 -55 -52 -53 ... and w'_n is empty if 4 divides n and $w'_n = (3-n, 4-n)$ otherwise. First patchworks and flip sequences are :

[-26 11 10 -21 -20 -23 -22 -2 -3 -18 -19 13 12-15-14 8 9 5 4 24 25 7 6 -17 -16 1]
[-28 11 10 -21 -20 -23 -22 -2 -3 -18 -19 13 12-15-14 8 9 5 4-25-24 26 27 7 6 -17 -16 1]
[-30 11 10 -21 -20 -23 -22 -2 -3 -18 -19-26-27-24-25 13 12-15-14 8 9 5 4 28 29 7 6 -17 -16 1]
[-32 11 10 -21 -20 -23 -22 -2 -3 -18 -19-26-27-24-25 13 12-15-14 8 9 5 4-29-28 30 31 7 6 -17 -16 1]
[-34 11 10 -21 -20 -23 -22 -2 -3 -18 -19-26-27-24-25-30-31-28-29 13 12-15-14 8 9 5 4 32 33 7 6 -17 -16 1]
[-36 11 10 -21 -20 -23 -22 -2 -3 -18 -19-26-27-24-25-30-31-28-29 13 12-15-14 8 9 5 4-33-32 34 35 7 6 -17 -16 1]
(26 18 10 14 11 3 24 11 16 8 19 5 13 25 8 21 5 18 3 15 26 15 3 18 21 8 25 11 13 7 19 8 16 11 24 3 11 16 14 20)
(28 20 10 16 13 3 26 11 18 2 10 21 5 15 27 8 21 5 18 3 15 28 17 3 20 23 8 27 13 15 7 21 10 2 18 11 26 3 13 18 16 2 22)
(30 22 14 18 15 3 28 11 16 21 23 10 21 5 15 29 10 23 5 20 26 24 3 15 30 19 3 22 25 8 29 15 17 7 23 12 20 11 28 3 11 15 20 18 11 24)
(32 24 14 20 17 3 30 11 18 2 23 25 12 23 5 17 31 10 23 5 20 26 24 3 15 32 21 3 24 27 8 31 17 19 7 25 14 2 22 11 30 3 13 17 22 20 13 2 26)
(34 26 18 22 19 3 32 11 16 25 27 23 25 12 23 5 17 33 12 25 5 22 28 26 30 28 3 15 34 23 3 26 29 8 33 19 21 7 27 16 24 11 32 3 11 15 19 24 22 15 11 28)

(36 28 18 24 21 3 34 11 18 2 27 29 25 27 14 25 5 19 35 12 25 5 22 28 26 30 28 3 15 36 25 3 28 31 8 35 21 23 7 29 18 2 26 11 34 3 13 17 21 26 24 17 13 2 30)
(40 32 22 28 25 3 38 11 18 2 31 33 29 31 27 29 16 27 5 21 39 14 27 5 24 30 28 32 30 34 32 3 15 40 29 3 32 35 8 39 25 27 7 33 22 2 30 11 38 3 13 17 21 25 30 28 21 17 13 2 34)
(44 36 26 32 29 3 42 11 18 2 35 37 33 35 31 33 29 31 18 29 5 23 43 16 29 5 26 32 30 34 32 36 34 38 36 3 15 44 33 3 36 39 8 43 29 31 7 37 26 2 34 11 42 3 13 17 21 25 29 34 32 25 21 17 13 2 38)
(48 40 30 36 33 3 46 11 18 2 39 41 37 39 35 37 33 35 31 33 20 31 5 25 47 18 31 5 28 34 32 36 34 38 36 40 38 42 40 3 15 48 37 3 40 43 8 47 33 35 7 41 30 2 38 11 46 3 13 17 21 25 29 33 38 36 29 25 21 17 13 2 42)
This provides even fortuitous sequences for all $n \geq 26$.

Double fortuitous sequences

For odd n a fortuitous sequence holds 3 f_n and may be of the form $(ns)^3$. As well, an even fortuitous sequence holds 2 f_n and may be of the form $(ns)^2$. Then its length $3n/2+1$ is even and $n \equiv 2 \pmod 4$.

An even fortuitous sequence is double whenever its patchwork verifies $f^{-1} = -f$. With this extra rule depth first search for even fortuitous sequences finds all double fortuitous sequences for $n = 30$(in 0.75s), $n = 34$(in 16.6s) and $n = 38$(in 9m). Some patchworks for 30 have similarities with some patchworks for 34, such as $f(2) = 8$. When adding this condition, computation is much faster. So we look for other similarities between patchworks for 30, 34 and 38 and add these conditions. Etc. At last we impose $f(2) = 8$, $f(n/2+8) = 4$, $f(6) = n/2$, $f(n/2+1) = n/2+3$ and $f(n/2+5) = n-1$. So we get double fortuitous sequences for $n = 30$, 34, 38, 42 and 46 in 0.036s.

E.g. double fortuitous sequences
$(30\ 22\ 14\ 7\ 3\ 24\ 7\ 27\ 13\ 15\ 10\ 29\ 15\ 20\ 18\ 5\ 23\ 14\ 20\ 13\ 28\ 3\ 19)^2$
$(34\ 26\ 16\ 7\ 3\ 28\ 9\ 31\ 13\ 17\ 2\ 12\ 33\ 17\ 24\ 2\ 22\ 5\ 27\ 18\ 2\ 24\ 15\ 32\ 3\ 21)^2$
$(38\ 30\ 18\ 7\ 3\ 32\ 7\ 35\ 15\ 17\ 15\ 19\ 14\ 37\ 19\ 26\ 28\ 26\ 22\ 9\ 31\ 18\ 26\ 28\ 26\ 17\ 36\ 3\ 23)^2$
$(42\ 34\ 20\ 7\ 3\ 36\ 9\ 39\ 15\ 17\ 15\ 21\ 2\ 16\ 41\ 21\ 30\ 32\ 30\ 2\ 26\ 9\ 35\ 22\ 2\ 30\ 32\ 30\ 19\ 40\ 3\ 25)^2$
$(46\ 38\ 22\ 7\ 3\ 40\ 14\ 16\ 12\ 14\ 11\ 43\ 14\ 18\ 21\ 23\ 14\ 45\ 27\ 35\ 31\ 28\ 26\ 13\ 39\ 22\ 32\ 35\ 33\ 37\ 35\ 21\ 44\ 3\ 27)^2$
have patchworks :
[-30 8 9-23-22 15 14 -2 -3-26-27-24-25 -7 -6 18 19-16-17 29 28 5 4 12 13 10 11-21-20 1]
[-34 8 9-25-24 17 16 -2 -3-28-29 31 30-26-27 -7 -6 20 21-18-19 33 32 5 4 14 15 10 11-13-12-23-22 1]
[-38 8 9-27-26 19 18 -2 -3-34-35-30-31-28-29-32-33 -7-6 22 23-20-21 37 36 5 4 14 15 12 13 16 17 10 11-25-24 1]
[-42 8 9-29-28 21 20 -2 -3-36-37 39 38-32-33-30-31-34-35 -7 -6 24 25-22-23 41 40 5 4 16 17 14 15 18 19 10 11 -13 -12 -27 -26 1]
[-46 8 9-31-30 23 22 -2 -3-42-43-40-41-37-36-39-38-33-32-35-34 -7 -6 26 27-24-25 45 44 5 4 19 18 21 20 15 14 17 16 12 13 10 11-29-28 1]

Palindromic even fortuitous sequences

An even fortuitous sequence is of the form $ns_1ns_2 = n\tilde{s}_2n\tilde{s}_1$ whenever $f = -f_n \circ f \circ f_n$. When adding this rule as well as $f(2) = n/2-2$, $f(6) = -2$, $f(10) = 3-n$ and $f(n-3) = 8$ for $n \neq 34$, depth first search for even fortuitous sequences finds readily even palindromic fortuitous sequences of the form $nw_nn\tilde{w}_n$ with
w_{30}=(24 14 12 20 15 3 28 11 16 14 20 14 23 5 17 29 8 23 7 20 3 17)
w_{34}=(28 18 12 16 24 15 3 32 15 20 14 24 18 27 5 21 33 8 23 27 29 9 22 3 19)
w_{38}=(32 22 20 28 21 23 21 3 36 11 20 18 24 18 29 31 29 5 17 37 12 31 7 26 28 26 3 21)
w_{42}=(36 26 24 32 25 2 27 23 3 40 11 22 20 26 20 31 35 2 33 5 17 41 14 35 7 28 32 2 30 3 23)
w_{46}=(40 30 28 36 31 15 7 3 44 27 35 39 35 15 20 18 24 18 27 29 37 13 25 45 12 39 15 36 34 23 19 30 3 25)
and their patchworks :
[-30 13 12-23-22 -2 -3 -6 -7-27-26-11-10 16 17-14-15 21 20 5 4 24 25 28 29 9 8-19-18 1]
[-34 15 14-25-24 -2 -3 -8 -9-31-30 -6 -7-22-23 17 16-19-18 12 13 28 29 5 4 26 27 32 33 11 10-21-20 1]
[-38 17 16-31-30 -2 -3 -6-7-35-34-15-14-11-10-13-12 20 21-18-19 27 26 29 28 25 24 5 4 32 33 36 37 9 8-23-22 1]
[-42 19 18-35-34 -2 -3 -6 -7-39-38-17-16-11-10 14 15-13-12 22 23-20-21 31 30-28-29 33 32 27 26 5 4 36 37 40 41 9 8-25-24 1]
[-46 21 20-39-38 -2 -3 -6 -7-43-42-11-10-13-12-29-28-33-32-17-16 24 25-22-23 31 30 15 14 19 18 35 34 37 36 5 4 40 41 44 45 9 8-27-26 1]

An even fortuitous sequence is of the form $ns_1ns_2 = n\tilde{s}_1n\tilde{s}_2$ whenever $f = f_n \circ f^{-1} \circ f_n$. When adding this rule as well as $f(2) = 7$, $f(4) = 11$, $f(7-n) = 13$, $f((n/2 \bmod 4)+7) = -3$ and $f((n/2 \bmod 4)+9) = 14$ depth first search for even fortuitous sequences finds readily even palindromic fortuitous sequences. After rotation these sequences take the form $(n{-}1\ w_n\ n{-}1\ \tilde{w}_n)$.
w_{34}=(24 12 20 27 9 23 32 26 30 17 25 34 27 17 6 8 30 9 20 32 6 27 12 19 11)
w_{38}=(28 16 2 24 31 11 27 36 30 34 19 27 38 31 19 8 10 34 11 2 22 36 8 2 31 12 21 13)
w_{42}=(28 12 24 35 31 27 9 23 40 34 38 25 33 42 35 25 14 16 12 14 10 12 38 9 20 40 10 35 12 23 15)
w_{46}=(32 16 2 28 39 35 31 11 27 44 38 42 27 35 46 39 27 16 18 14 16 12 14 42 11 2 22 44 12 2 39 12 25 17)
[-34 7 6 11 10-23-22 -3 -2 14 15-29-28 -5 -4-17-16-19-18 24 25 -8 -9 31 30-12-13 33 32-21-20-27-26 1]
[-38 7 6 11 10 24 25-23-22 -3 -2 14 15 32 33-31-30 -5-4-17-16-19-18 26 27-8-9 35 34-12-13 37 36-21-20-29-28 1]
[-42 7 6 11 10-31-30 -3 -2 14 15-37-36 -5 -4-25-24-27-26-21-20-23-22-17-16-19-18 32 33 -8 -9 39 38-12-13 41 40-29-28-35-34 1]
[-46 7 6 11 10 32 33-31-30 -3 -2 14 15 40 41-39-38 -5 -4-25-24-27-26-21-20-23-22-17-16-19-18 34 35 -8 -9 43 42-12-13 45 44-29-28-37-36 1]

An even fortuitous sequence is of the form $(ns)^2 = (n\tilde{s})^2$ whenever $-f = f^{-1} = f_n \circ f \circ f_n$. When adding this rule depth first search for even fortuitous sequences finds readily double palindromic fortuitous sequences
$(34w_{34}33\tilde{w}_{34})^2$ with w_{34}=(14 5 3 30 14 10 5 3 18 6 21 10)
$(42w_{42}41\tilde{w}_{42})^2$ with w_{42}=(31 27 4 35 33 14 40 20 34 12 23 6 20 39 34) and their patchworks :
[-34 20 21-26-27 17 16-30-31-12-13 10 11 32 33 -7 -6 29 28 -2 -3-24-25 22 23 4 5-19-18 8 9-14-15 1]
[-42 11 10-19-18-29-28 16 17 -3 -2-23-22-37-36 -8 -9 5 4-31-30 13 12-39-38 34 35 7 6 21 20 41 40-26-27 15 14 25 24-33-32 1]

Generalized odd fortuitous sequences

For odd n fortuitous sequences may be generalized by replacing the stack of 2-clans placed on singleton 1, by a patchwork of 2-clans and of 2-blocks (placed on singleton 1). So we define an "generalized odd fortuitous sequence" as a flip sequence of length $\frac{3n+3}{2}$ of the form $S = ns_1ns_2ns_3$ where s_1, s_2 and s_3 are sequences of positive integers lower than n, such that $f_S = -I_n$ and stack $f = -f_{ns_1}$ obtained at the end of s_1, when applying S from $-I_n$ or equivalently s_1 from $-f_n$, is a "patchwork" of $\frac{n-1}{2}$ clans or blocks of 2 pancakes, placed on pancake 1.

So no pancake of this patchwork has potential 3 (or -3) and every flip of s_1 increases potential by 4, and breaks an anti-adjacency. As well every flip of s_2 and of s_3 creates an adjacency. We may notice that $h = f_n \circ f^{-1} \circ f_n$ is the patchwork of generalized fortuitous sequence $n\tilde{s}_1n\tilde{s}_3n\tilde{s}_2$, and $\tilde{s}_1$, $\tilde{s}_3$, s_1, and s_2 are the only flip sequences of their lengths adding an adjacency at each step when starting respectively from stacks $-f$, f^{-1}, $-f_n \circ f^{-1}$ and $f_n \circ f$. So patchwork f characterizes s_1, since its length is the number of missing anti-adjacencies in f.

Sequence s_1 derives from f as $\tilde{s}_1$ derives from h. Hence $f = h$ if and only if $s_1 = \tilde{s}_1$, i.e. if s_1 is a palindrome. E.g. for $n = 15$ or $n = 19$ there is only one generalized fortuitous sequence. Hence $f = h$ and s_1 is a palindrome and $s_3 = \tilde{s}_2$. Hence $s_3ns_1ns_2$ is a palindrome.

Any fortuitous sequence is a generalized fortuitous sequence. Hence there are more generalized fortuitous sequences than fortuitous sequences and hence more chance to find generalized fortuitous sequences than fortuitous sequences. E.g. for $n = 19$, there is no fortuitous sequence, but there is a generalized fortuitous sequence. As well for $n = 25$, no fortuitous sequence is known, but a generalized fortuitous sequence is easily found. Patchworks of generalized fortuitous sequences are permutations of $\{-n, 1{-}n, \ldots -1, 0, 1, 2, \ldots n\}$ verifying following conditions:

$*\forall i,\ f(-i) = -f(i)$	f is a signed permutation.
$*f(n) = 1$	Smallest pancake is at bottom right-side up.
$*\forall i,\ f(i - d(i)) = f(i) + d(f(i))$	f is made of clans and blocks of 2 pancakes.
$*\forall i,\ 2f(i) \neq f(i-1) + f(i+1)$	Clans and blocks of 2 pancakes do not form larger clans or blocks.

We perform a depth first search to find all generalized fortuitous sequences. At start patchwork f and stacks $-f$, f^{-1}, $-f_n \circ f^{-1}$ and $f_n \circ f$ have most of their values unknown. To make sure a flip on one of the four stacks creates an adjacency, we may have to define some values of f. We choose the stack which minimizes the number of unknown values of f to define. E.g. a stack of the form $[x \ldots y \ldots]$ is best choice,

if values x and $y = 1 - x$ are known, while a stack with an unknown top pancake is worst choice. Once the stack is chosen we try any possible flip on it and choose again a stack. When the top of a stack is 1 or $-n$ we may force the stack to be I_n or f_n respectively.

This depth first search finds generalized fortuitous sequences for $n = 15$, 19 and 25(in 6m 6s). We may look for similarities between a solution for 15 and a solution for 19 and impose them. This gives readily solutions for 15, 19, 23 and 27. Then looking for other similarities between those solutions, we impose, when $n \equiv 3 \pmod 4$ and $n \geq 15$, that $f =$ [-8 -9 w_n -12 -13 -2 -3 1-n -n 5 4 7 6 -10 -11 1], where w_n is a terminal segment of length $n - 15$, which is multiple of 4, of infinite sequence :
... -39 -38 -41 -40|-35 -34 -37 -36|-31 -30 -33 -32|-27 -26 -29 -28|-23 -22 -25 -24|-19 -18 -21 -20|-15 -14 -17 -16
Fortuitous sequences are of the form (n u_n n−1 $\tilde{u}_n$) since $f = h$. First patchworks and u_n are :
[-8 -9 -12 -13 -2 -3 -14 -15 5 4 7 6 -10 -11 1]
[-8 -9 -15 -14 -17 -16 -12 -13 -2 -3 -18 -19 5 4 7 6 -10 -11 1]
[-8 -9 -19 -18 -21 -20 -15 -14 -17 -16 -12 -13 -2 -3 -22 -23 5 4 7 6 -10 -11 1]
[-8 -9 -23 -22 -25 -24 -19 -18 -21 -20 -15 -14 -17 -16 -12 -13 -2 -3 -26 -27 5 4 7 6 -10 -11 1]
[-8 -9 -27 -26 -29 -28 -23 -22 -25 -24 -19 -18 -21 -20 -15 -14 -17 -16 -12 -13 -2 -3 -30 -31 5 4 7 6 -10 -11 1]
u_{15} = (10 4 6 14 6 4 10 15 10 4 6)
u_{19} = (14 7 4 10 18 6 4 10 19 14 4 9 11 8)
u_{23} = (18 11 7 4 14 22 6 4 10 23 18 4 13 15 11 13 10)
u_{27} = (22 15 11 7 4 18 26 6 4 10 27 22 4 17 19 15 17 13 15 12)
u_{31} = (26 19 15 11 7 4 22 30 6 4 10 31 26 4 21 23 19 21 17 19 15 17 14)

As well, when $n \equiv 1 \pmod 4$ and $n \geq 25$, we impose
$f =$ [-8 -9 n-3 n-2 n-7 n-6 -12 -13 -2 -3 w_n 5 4 1-n -n w'_n 5-n 4-n 7 6 -10 -11 1], where w_n is an terminal segment of even length $\frac{n-21}{2}$ of infinite sequence: ... 49 48|45 44|41 40|37 36|33 32|29 28|25 24|21 20|17 16
and w'_n is an initial segment of even length $\frac{n-21}{2}$ of infinite sequence: 15 14|19 18|23 22|27 26|31 30|35 34 ...
E.g. patchworks and flip sequences for 25, 29 and 33 are :
[-8 -9 22 23 18 19 -12 -13 -2 -3 17 16 5 4 -24 -25 15 14 -20 -21 7 6 -10 -11 1]
[-8 -9 26 27 22 23 -12 -13 -2 -3 21 20 17 16 5 4 -28 -29 15 14 19 18 -24 -25 7 6 -10 -11 1]
[-8 -9 30 31 26 27 -12 -13 -2 -3 25 24 21 20 17 16 5 4 -32 -33 15 14 19 18 23 22 -28 -29 7 6 -10 -11 1]
(25 10 20 4 13 17 10 24 12 15 9 4 11 17 20 25 16 4 12 24 10 4 20 6 13 9 25 18 15 7 4 10 22 10 24 6 20 4 10)
(29 14 24 4 13 17 10 28 16 8 10 19 11 6 13 21 24 29 20 6 16 28 10 4 22 8 24 22 13 9 29 22 19 11 4 14 6 26 10 28 10 6 24 4 10)
(33 18 28 4 13 17 10 32 20 12 14 10 12 23 13 8 15 25 28 33 24 8 20 32 10 4 24 10 26 24 28 26 13 9 33 26 23 15 4 18 6 10 30 10 32 14 10 6 28 4 10)
This provides odd generalized fortuitous sequences for all $n \geq 23$.

Results

We have produced for every integer $n \in \{15, 19, 23\}$ or $n \geq 25$, an explicit patchwork yielding an even fortuitous sequence or a generalized odd fortuitous sequence (according to parity of n), proving thereby that for all these numbers $g(-I_n) = \lceil \frac{3}{2}n + 1 \rceil$. We can achieve the same result in another way. We have fortuitous sequences for $n \in \{15, 23, 31, 29, 33, 37, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48\}$ produced as palindromic fortuitous sequences when $n \equiv 3 \pmod 4$ or as triple fortuitous sequences for odd $n \geq 29$ or as even fortuitous sequences for even n. Each of these sequences may be checked. We may then complete this set of fortuitous sequences by combining every sequence with the sequence for 15 or 26 according to its parity. So we get fortuitous sequences for all $n \geq 26$. With the two generalized fortuitous sequences for 19 and 25 result is achieved. Of course we may also consider fortuitous sequences $(2\ 1)^2$ and $(3\ 2)^3$.

All fortuitous sequences shown in this paper may be computed by program in C gnu of annex. To get computational times given here, program was compiled with gcc 4.8.3 and run on an AMD FX-4130. E.g. `triple(37)` finds and prints 24 triple fortuitous sequences in 10.5s for n=37 if `sm`=0 but `triple(39)` finds and prints 5 triple palindromic fortuitous sequences at once for n=39 if `sm` $\neq$ 0. As well `palin(23)` finds and prints 4 palindromic fortuitous sequences in 0.008s for n=23 and `simple(n)` looks for one (if `ALL`=0) or all generalized odd fortuitous sequences or even fortuitous sequences according to parity of n. If `sm`=1 it adds condition $f = -f^{-1}$ and computes only double sequences. If `sm`=2 it adds condition $f = f_n \circ -f \circ f_n$ and computes only palindromic sequences centered on n−1. If `sm`=4 it adds condition $f = f_n \circ f^{-1} \circ f_n$

and computes only palindromic sequences centered on n. If `sm`=3 it adds the 3 conditions and computes only palindromic double sequences. If `DIR` $\neq 0$ and `sm`=0 it adds conditions such as $f(1) = -8$ etc. and produces at once an imposed fortuitous sequence. Program combines also small given fortuitous sequences to produce and print a fortuitous sequence for every n from 26 to 199, and checks that produced sequences are fortuitous. Program is on `http://mapage.noos.fr/laurent.pierre/concat.c`.

Bibliography

[1] William H. Gates, Christos H. Papadimitriou, Bounds for sorting by prefix reversal, Discrete Mathematics, Volume 27, Issue 1, 1979, Pages 47-57, ISSN 0012-365X

[2] David S. Cohen, Manuel Blum, On the problem of sorting burnt pancakes, Discrete Applied Mathematics, Volume 61, Issue 2, 28 July 1995, Pages 105-120, ISSN 0166-218X

[3] Mohammad H. Heydari and I. Hal Sudborough. 1997. On the diameter of the pancake network. J. Algorithms 25, 1 (October 1997), 67-94

[4] Josef Cibulka, On average and highest number of flips in pancake sorting, Theoretical Computer Science, Volume 412, Issues 8-10, 4 March 2011, Pages 822-834, ISSN 0304-3975

Annex : program in C gnu

```
#include<stdio.h>
#include<stdlib.h>
#define mmax 100
#define DIR 1
#define ALL 0
typedef int stack[2*mmax-1], seq[3*mmax];
void prnt(int *t, int n) { while(n--) printf("%3d",*t++); printf("\n"); }
int aff=1, sm=0;
void pseq(seq q) { if(~aff) prnt(q,(*q+1)*3/2); }
void prsq(stack s, seq q) { if(aff) prnt(s,*q); pseq(q); }
void flip(stack s, int l) {while(l>0) {int x=s[--l]; s[l--]=-*s, *s++=-x;}}
void copy(stack s, stack t, int n) { for(;n--;) t[n]=s[n]; }       // t=s
void minv(stack s, stack t, int n)                                 // t=-In/s
{ for(;n--;) if(s[n]<0) t[~s[n]]=n+1; else t[s[n]-1]=~n; }
int isIn(stack s, int n) { for(;n && s[n-1]==n;n--) ; return !n; } // s==In
void appl(stack s, seq q)
{ int n=*q, i;
  for(i=n;i--;) s[i]=~i;
  if(n&3) q[n&1?n/2+1:n*3/4+1]=n;
  do flip(s,*q++); while(*q!=n);
}
void chkseq(seq q)
{ int n=*q, i; stack s;
  for(i=n;i--;) s[i]=~i;
  for(i=3*(n+1)/2;i--;) flip(s,q[i]);
  if(!isIn(s,n)) printf("wrong sequence\n"), exit(1);
}
int*sort(stack s, seq q, int n)
{ int i=n, x; for(;;) for(flip(s,*q++=i),i=1,x=1-*s;s[i]!=x;) if(++i==n) return q; }
int stsq(stack s, seq q, int n)
{ stack t;
  minv(s,t,n); int *r=sort(t,q,n),      c=1&n;    if(   !isIn(t,n)) return 0;
  copy(s,t,n); do   r=sort(t,r,n); while(c--);    return isIn(t,n) && r-q==(n+1)*3/2;
}
void compose(seq a, seq b, seq c)
{ stack s, t;   appl(s,a), appl(t,b);  // t[]={-l-2,-l-3,y,y-1,...}
```

```
  int d=*a&1?2:1, i, l=-*t-2, h=*a+~d, x;
  for(i=0;i<*a;i++) x=s[i],s[i  ]=x+=x>0  ?l:     -l  ;  // |x|+=l
  for(i=d;i<*b;i++) x=t[i],s[i+h]=x+=x>-*t?h:x<*t?-h:0;  // |x|+=h if |x|>l+2
  if(!stsq(s,c,i+h)) printf("compose failed\n"),exit(1);
}
#define N(i) (i>0?i-n-1:i+n+1)
#define I(x,y) x=(i>0?(i-n)^1:-((n-i)^1))+n, y=j>0?j^1:-(-j^1)
void trip(int*f, int*h, int n) //-8 -9-12-13 -2 -3-14-15  5  4  7  6-10-11  1
{ void set(int i, int j)
  { int d=2*(1&(i^i>>10))-1;
    if(f[i]!=h[j] || f[i] || (f[j] && h[i] && f[j]!=h[i])
                  || f[i-d]==j+d || f[1]>0) return;
    h[f[i]=j]=i, h[f[-i]=-j]=-i;
    if(!f[i+d]) set(i+d,j-d);
    else if(!f[j]&& h[i]) set(j,h[i]);
    else if( f[j]&&!h[i]) set(f[j],i);
    else if(sm && f[N(j)]!=N(i)) set(N(j),N(i));
    else trip(f,h,n);
    f[i]=f[-i]=h[j]=h[-j]=0;
  }
  int i, j, k=0, c=0; static seq q;
  for(i=n;i;i--) if(!f[i] && h[c=i] && h[k=h[i]]) return set(i,h[k]);
  if(!c) if(stsq(f+1,q,n)) prsq(f+1,q);
  if(k) for(i=f[k],j=n;j;j--) if(h[j]==f[j]) set(i,(i^j)&1?-j:j);
}
void triple(int n)
{ int i=2*n+1, s[i], t[i], *f=s+n, *h=t+n;
  for(;i--;) s[i]=t[i]=0;
  h[f[n]=1]=n, h[f[-n]=-1]=-n;
  void E(int i,int j) { for(;!f[i];I(i,j)) h[f[i]=j]=i, h[f[-i]=-j]=-i;}
  if(DIR && !sm) E(-1,8),E(4,12-(i=n|2)),E(-5,2),E(-8,n),E(9,5),E(11,7-n),E(i-15,n-5);
  trip(f,h,n);
}
void palin(int i)
{ static seq q; static stack s, t; static int k=0, j, m, n;
  if(!k) for(j=n=i,m=n/2+1;j--;) s[j]=~j;
  flip(s,q[k++]=i);
  if(n&2 && k<m && (j=m-k)<=k) { i=j<k?q[j]:n-1; if(s[i]==s[i-1]-1) palin(i); } else
  if(k<m) for(i=2;i<n;i+=2)    {                 if(s[i]==s[i-1]-1) palin(i); } else
  if(copy(s,t,n), *s<0 && sort(t,q+m,n)==q+2*m && sort(t,q+2*m,n)==q+3*m) pseq(q);
  flip(s,q[--k]);
}
void simple(int n)
{ seq q={n}, r={n}; static stack s, S, t, T;
  int u[8*mmax], *f=u+2*mmax, *h=f+4*mmax, kK=(n+1)*3/2, k=1, K=kK-k, l=k, L=K, i;
  void dfs()
  { void set(i,j)
    { int I(I,J);
      if(i && j && f[2*i-I]!=2*j-J && f[2*I-i]!=2*J-j)
      h[f[i]=j]=i, h[f[-i]=-j]=-i, h[f[I]=J]=I, h[f[-I]=-J]=-I,
      (sm&2 && !f[N(i)]? set(N(i),-N(j)) :sm&1 && !f[j]? set(j,-i)
      :sm&4 && !f[N(j)]? set(N(j), N(i)) :dfs()),
      f[i]=h[j]=f[-i]=h[-j]=f[I]=h[J]=f[-I]=h[-J]=0;
```

```
    }
    int x=f[*s], X=f[*S], y=x?h[1-x]:0, Y=X?h[1-X]:0,
        a=h[*t], A=h[*T], b=a?f[1-a]:0, B=A?f[1-A]:0, i;
    if(k==K) { if(~n&1) q[k]=n, prsq(f+1,q), n=ALL?n:1; return; }
    if(l==L) { if(~n&1) r[l]=n, prsq(h+1,r), n=ALL?n:1; return; }
    if(K+l==kK && k+l==L && X==-n && a==1 && n&1)
       return r[L]=n, copy(q,r+l,k), prsq(f+1,r), n=ALL?n:1, chkseq(r);
    #define SET(a,b) if(f[a]!=b) { if(!f[a] && !h[b]) set(a,b); return; }
    if(x==1)  for(i=n;--i;) SET(s[i],i+1)
    if(X==-n) for(i=n;--i;) SET(S[i],i-n)
    if(a==1)  for(i=n;--i;) SET(i+1,t[i])
    if(A==-n) for(i=n;--i;) SET(i-n,T[i])
    i=n; if(y) {for(;--i;) if(s[i]==y) flip(s,q[k++]=i), dfs(), flip(s,q[--k]), i=1;}
    else if(Y) {for(;--i;) if(S[i]==Y) flip(S,q[K--]=i), dfs(), flip(S,q[++K]), i=1;}
    else if(b) {for(;--i;) if(t[i]==b) flip(t,r[l++]=i), dfs(), flip(t,r[--l]), i=1;}
    else if(B) {for(;--i;) if(T[i]==B) flip(T,r[L--]=i), dfs(), flip(T,r[++L]), i=1;}
    else if(X) {for(;--i;) if(!f[S[i]]) set(S[i],1-X);}
    else if(a) {for(;--i;) if(!h[t[i]]) set(1-a,t[i]);}
    else if(x) {for(;--i;) if(!f[s[i]]) set(s[i],1-x);}
    else if(A) {for(;--i;) if(!h[T[i]]) set(1-A,T[i]);}
    else        for(;--i;) if(!h[i]) set(*s,i), set(*s,-i);
  }
  void E(int i,int j) { for(;!f[i];I(i,j)) { h[f[i]=j]=i, h[f[-i]=-j]=-i;
    if(sm&1) E(j,-i); if(sm&2) E(N(i),-N(j)); if(sm&4) E(N(j),N(i)); } }
  for(i=n;i;) i--, h[s[i]=i-n]=h[t[i]=n-i]=f[S[i]=~i]=f[T[i]=i+1]=0;
  *f=*h=f[~n]=h[~n]=f[n+1]=h[n+1]=0, f[h[1]=n]=1, f[h[-1]=-n]=-1;
  if(~n&1) h[f[1]=-n]=1, f[h[n]=-1]=n;
  if(!DIR || sm&(sm-1)) ; else
  if(sm==1) E(2,8),E(6,n/2),E(n/2+8,4),E(n/2+1,n/2+3),E(n/2+5,n-1); else
  if(sm==2) E(2,n/2-2),E(6,-2),E(10,3-n),E(n-3,n!=34?8:10); else
  if(sm==4) E(2,7),E(4,11),E(n-7,-13),E(n/2%4+7,-3),E(n/2%4+9,14); else
  if(n&1) if(E(-1,8),E(1-n,11),E(n-3,6),n&2) for(E(n-5,4),E(7-n,n),E(9-n,3),E(11-n,13);
                  n+i>15;i-=4) E(13-n-i,16-i),E(15-n-i,14-i);
          else for(E(3,n-3),E(5,n-7),E(-7,12),E(-9,2), E(n/2+1,5),E(-n/2-4,n),E(5-n,n-4);
                  i+24<n;i+=4) E(n/2+i/2+6,14+i),E(n/2-i/2,16+i);
  else for(E(2,11),E(-4,21),E(-6,23),E(-8,2),E(-10,18), E(n-5,n-1),E(n-3,6),E(1-n,16);
          n+i>30;i-=4) E(i-12,26-i),E(i-14,24-i);
  dfs();
}
seq ss[2*mmax]={{2,1},     {3,2},       {31,22,16,18,30,16,22,24,20,22,16,6,18,26,12,18},
{29,24,4,12,28,14,6,24,16,4,20,6,26,4,16},  {33,18,8,30,8,2,20,4,14,8,4,24,8,26,32,4,18},
{37,22,12,2,34,8,20,4,16,2,10,4,26,2,8,28,36,6,20},   {23,14,4,6,22,10,8,12,10,14,12,18},
{30,22,14,7,3,24,7,27,13,15,10,29,15,20,18,5,23,14,20,13,28,3,19}, {15,10,4,6,14,6,4,10},
{42,34,20,7,3,36,9,39,15,17,15,21,2,16,41,21,30,32,30,2,26,9,35,22,2,30,32,30,19,40,3,
25}, {38,30,18,7,3,32,7,35,15,17,15,19,14,37,19,26,28,26,22,9,31,18,26,28,26,17,36,3,23},
{46,38,22,7,3,40,14,16,12,14,11,43,14,18,21,23,14,45,27,35,31,28,26,13,39,22,32,35,33,37,
35,21,44,3,27},  {34,26,16,7,3,28,9,31,13,17,2,12,33,17,24,2,22,5,27,18,2,24,15,32,3,21},
{26,18,10,14,11,3,24,11,16,8,19,5,13,25,8,21,5,18,3,15},
{28,20,10,16,13,3,26,11,18,2,10,21,5,15,27,8,21,5,18,3,15,28},
{32,21,3,24,27,8,31,17,19,7,25,14,2,22,11,30,3,13,17,22,20,13,2,26,32},
{48,37,3,40,43,8,47,33,35,7,41,30,2,38,11,46,3,13,17,21,25,29,33,38,36,29,25,21,17,13,2,
42,48},     {36,25,3,28,31,8,35,21,23,7,29,18,2,26,11,34,3,13,17,21,26,24,17,13,2,30,36},
{44,33,3,36,39,8,43,29,31,7,37,26,2,34,11,42,3,13,17,21,25,29,34,32,25,21,17,13,2,38,44},
```

```
{40,29,3,32,35,8,39,25,27,7,33,22,2,30,11,38,3,13,17,21,25,30,28,21,17,13,2,34,40}, {0}};
int main()
{ int i, j, h=0, *s[2*mmax];
  for(i=2*mmax;i--;) s[i]=0;
  for(;i=*ss[h],i ;) s[i]=ss[h++],  compose(s[2+i%2],s[i],s[i]);
  for(i=27;i<2*mmax;i+=2) if(!s[i]) compose(s[i-12],s[15],s[i]=ss[h++]);
  for(i=50;i<2*mmax;i+=2)           compose(s[i-24],s[26],s[i]=ss[h++]);
  for(i= 0;i<2*mmax;i++ ) if( s[i]) pseq(s[i]);
  palin(15), palin(23), triple(15), simple(15), simple(19), simple(25);
  for(i=15;i<(DIR||sm?48:38);i+=2) triple(i);
  for(j=4;j--;) for(aff=-1;aff<1;aff++) for(i=15;i<60;i++ ) if(i%4==j) simple(i);
  while(++sm<5) for(prnt(&sm,1),i=sm>2?34:30;i<50;i+=4<<(sm==3)) simple(i);
  return 0;
}
```